

Datasheet for ABIN1404049

anti-C19orf28 antibody (Biotin)



Go to Product page

()	V		rV	ĺ	9	V	V
'	\mathcal{I}	٧V	<u> </u>	v	1	$\overline{}$	٧	٧

Quantity:	100 μL
Target:	C19orf28
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This C19orf28 antibody is conjugated to Biotin
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MFSD12/C19orf28	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Purified by Protein A.	

Target Details

Target:

C19orf28

. a. get.	- 130.1. 2 0
Alternative Name:	C19orf28 (C19orf28 Products)
Background:	Synonyms: Chromosome 19 open reading frame 28, Hypothetical protein LOC126321,
	MGC20700, PP3501, Uncharacterized MFS type transporter C19orf28, MFS12_HUMAN.
	Background: C19orf28, also known as PP3501, is a multi-pass membrane protein that belongs

to the major facilitator superfamily. The gene encoding C19orf28 localizes to chromosome 19

and, due to alternative splicing events, C19orf28 exists as two isoforms. Consisting of around 63 million bases with over 1,400 genes, chromosome 19 makes up over 2 % of human genomic DNA. Chromosome 19 includes a diversity of interesting genes and is recognized for having the greatest gene density of the human chromosomes. It is the genetic home for a number of immunoglobulin superfamily members including the killer cell and leukocyte Ig-like receptors, a number of ICAMs, the CEACAM and PSG family, and Fc receptors. Key genes for eye color and hair color also map to chromosome 19. Peutz-Jeghers syndrome, spinocerebellar ataxia type 6, the stroke disorder CADASIL, hypercholesterolemia and insulin-dependent diabetes have been linked to chromosome 19. Translocations with chromosome 19 and chromosome 14 can be seen in some lymphoproliferative disorders and typically involve the proto-oncogene BCL3. The C19orf28 gene product has been provisionally designated C19orf28 pending further characterization.

Application Details

Application Notes:	IHC-P 1:200-400
Restrictions:	For Research Use only
Handling	

Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months