antibodies.-online.com

## Datasheet for ABIN1404239 anti-NHEDC2 antibody (Alexa Fluor 555)



| Overview          |  |
|-------------------|--|
| Quantity:         | 100 µL   |
| Target:           | NHEDC2   |
| Reactivity:       | Human, Mouse, Rat  |
| Host:             | Rabbit   |
| Clonality:        | Polyclonal   |
| Conjugate:        | This NHEDC2 antibody is conjugated to Alexa Fluor 555  |
| Application:      | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))  |
| Product Details   |  |
| Immunogen:        | KLH conjugated synthetic peptide derived from human NHEDC2   |
| Isotype:          | lgG  |
| Cross-Reactivity: | Human, Mouse, Rat  |
| Purification:     | Purified by Protein A.   |
| Target Details    |  |
| Target:           | NHEDC2   |
| Alternative Name: | NHEDC2 (NHEDC2 Products)   |
| Background:       | Synonyms: NHA2, Mitochondrial Na+/H+ exchanger NHA2, Mitochondrial sodium/hydrogen exchanger NHA2, Na+/H+ exchanger like domain containing protein 2, Na+/H+ exchanger |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1404239 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

exchanger like domain containing protein 2, SL9B2\_HUMAN.

|                     | Background: Na+/H+ exchangers (NHEs) catalyze the transport of Na+ in exchange for H+                              |
|---------------------|--|
|                     | across membranes in organisms and are required for numerous physiological processes.                               |
|                     | NHEDC2 (Na+/H+ exchanger-like domain-containing protein 2), also known as NHA2, is a 537                           |
|                     | amino acid mitochondrial protein. NHEDC2 is involved in organelle volume homeostasis by                            |
|                     | catalyzing the exchange of protons for Na+ and Li+ across the inner mitochondrial membrane.                        |
|                     | Found in red blood cells, NHEDC2 is required for bone resorption activity and osteoclast                           |
|                     | differentiation. As a multi-pass membrane protein, NHEDC2 is expressed as two isoforms                             |
|                     | produced by alternative splicing events.   |
| Gene ID:            | 133308   |
| Pathways:           | Proton Transport   |
| Application Details |  |
| Application Notes:  | IF(IHC-P) 1:50-200   |
| Restrictions:       | For Research Use only  |
| Handling            |  |
| Format:             | Liquid   |
| Concentration:      | 1 μg/μL  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.        |
| Preservative:       | ProClin  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:            | -20 °C   |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:        | 12 months  |