.-online.com antibodies

Datasheet for ABIN1404347 anti-UFD1L antibody (Alexa Fluor 555)



Overview	
Quantity:	100 µL
Target:	UFD1L
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UFD1L antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human UFD1L
Immunogen: Isotype:	KLH conjugated synthetic peptide derived from human UFD1L
Isotype:	IgG
Isotype: Cross-Reactivity:	lgG Human, Mouse, Rat
Isotype: Cross-Reactivity: Purification:	lgG Human, Mouse, Rat
Isotype: Cross-Reactivity: Purification: Target Details	IgG Human, Mouse, Rat Purified by Protein A.
Isotype: Cross-Reactivity: Purification: Target Details Target:	IgG Human, Mouse, Rat Purified by Protein A. UFD1L
Isotype: Cross-Reactivity: Purification: Target Details Target: Alternative Name:	IgG Human, Mouse, Rat Purified by Protein A. UFD1L UFD1L (UFD1L Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1404347 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

ubiquitin-dependent proteolytic pathway which degrades ubiquitin fusion proteins. This

	complex, also containing UFD1L, VCP and NPLOC4, binds ubiquitinated proteins and is required
	for the export of misfolded proteins from the ER to the cytoplasm for disposal. The NPLOC4-
	UFD1L-VCP complex also regulates spindle disassembly at the end of mitosis and is necessary
	for the formation of a closed nuclear envelope. UFD1L gene hemizygosity is the cause of some
	developmental defects including DiGeorge syndrome (DGS), velo-cardio-facial syndrome
	(VCFS) and Opitz G/BBB syndrome. UFD1L has 2 named isoforms produced by alternative
	splicing.
Gene ID:	7353
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months