antibodies -online.com





anti-FGF14 antibody (AbBy Fluor® 350)



Go to Product page

()	11/0	K\ /	iew
	\cup	ועוי	$I \cap VV$

Quantity:	100 μL	
Target:	FGF14	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This FGF14 antibody is conjugated to AbBy Fluor® 350	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FGF14
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	FGF14
Alternative Name:	FGF14 (FGF14 Products)
Background:	Synonyms: FHF 4, FHF-4, FHF4, Fibroblast growth factor 14, Fibroblast growth factor homologous factor 4, mFHF-41B, FGF14_HUMAN.
	Background: Fibroblast growth factor-1 (FGF-1), also designated acidic FGF, and fibroblast
	growth factor-2 (FGF-2), also designated basic FGF, are members of a family of growth factors

Target Details

that stimulate proliferation of cells of mesenchymal, epithe-lial and neuroectodermal origin. Additional members of the FGF family include the oncogenes FGF-3 (Int2) and FGF-4 (hst/Kaposi), FGF-5, FGF-6, FGF-7 (KGF), FGF-8 (AIGF), FGF-9 (GAF) and FGF-10FGF-23. Members of the FGF family share 30-55 % amino acid sequence identity and similar gene structure, and are capable of transforming cultured cells when overexpressed in transfected cells. Cellular receptors for FGFs are members of a second multigene family including four tyrosine kinases, designated Flg (FGFR-1), Bek (FGFR-L), TKF and FGFR-3.

Gene ID:

2259

Application Details

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
	00.00

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months