# antibodies -online.com







## anti-FGF14 antibody (Alexa Fluor 488)



$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Overview		
Quantity:	100 μL	
Target:	FGF14	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This FGF14 antibody is conjugated to Alexa Fluor 488	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Droduot Dotaile		

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human FGF14	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	

## Target Details

Target:	FGF14	
Alternative Name:	FGF14 (FGF14 Products)	
Background:	Synonyms: FHF 4, FHF-4, FHF4, Fibroblast growth factor 14, Fibroblast growth factor	
	homologous factor 4, mFHF-41B, FGF14_HUMAN.	
	Background: Fibroblast growth factor-1 (FGF-1), also designated acidic FGF, and fibroblast	
	growth factor-2 (FGF-2), also designated basic FGF, are members of a family of growth factors	

#### **Target Details**

that stimulate proliferation of cells of mesenchymal, epithe-lial and neuroectodermal origin. Additional members of the FGF family include the oncogenes FGF-3 (Int2) and FGF-4 (hst/Kaposi), FGF-5, FGF-6, FGF-7 (KGF), FGF-8 (AIGF), FGF-9 (GAF) and FGF-10FGF-23. Members of the FGF family share 30-55 % amino acid sequence identity and similar gene structure, and are capable of transforming cultured cells when overexpressed in transfected cells. Cellular receptors for FGFs are members of a second multigene family including four tyrosine kinases, designated Flg (FGFR-1), Bek (FGFR-L), TKF and FGFR-3.

Gene ID:

2259

### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 a 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should handled by trained staff only.	
Storage.	-20 °C	

handled by trained staff only.

Storage:

-20 °C

Storage Comment:

Store at -20 °C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date:

12 months