# antibodies -online.com





## Datasheet for ABIN1404577

# anti-C10orf93 antibody (Biotin)



#### Overview

Quantity:	100 μL
Target:	C10orf93 (TTC40)
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This C10orf93 antibody is conjugated to Biotin
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human C10orf93
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

### **Target Details**

Target:	C10orf93 (TTC40)
Alternative Name:	C10orf93 (TTC40 Products)
Background:	Synonyms: TTC40, C10orf92, C10orf93, C10orf123, C10orf124, bA288G11.4, bA288G11.5, bB137A17.2, bB137A17.3, Protein CFAP46, Cilia- and flagella-associated protein 46,
	Tetratricopeptide repeat protein 40, CFAP46
	Background: C10orf93, also known as C10orf124 or TPR repeat-containing protein C10orf93, is

a 1,530 amino acid protein that contains two TPR repeats and exists as three alternatively spliced isoforms. The gene encoding C10orf93 maps to human chromosome 10q26.3. Spanning nearly 135 million base pairs, chromosome 10 makes up approximately 4.5 % of total DNA in cells and encodes nearly 1,200 genes. Several protein-coding genes, including those that encode for chemokines, cadherins, excision repair proteins, early growth response factors (Egrs) and fibroblast growth receptors (FGFRs), are located on chromosome 10. Defects in some of the genes that map to chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss syndrome, Usher syndrome, nonsyndromatic deafness, Wolman?s syndrome, Cowden syndrome, multiple endocrine neoplasia type 2 and porphyria.

Gene ID:

54777

UniProt:

Q8IYW2

#### **Application Details**

Application Notes:

IHC-P 1:200-400

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months