-online.com antibodies

Datasheet for ABIN1404601 anti-SKIDA1 antibody (Biotin)



Overview	
Quantity:	100 μL
Target:	SKIDA1
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SKIDA1 antibody is conjugated to Biotin
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human C10orf140
lsotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.
Target Details	
Target:	SKIDA1
Alternative Name:	C10orf140 (SKIDA1 Products)
Background:	Synonyms: C10orf140, chromosome 10 open reading frame 140, DKFZp761J229, DLN 1, DLN1_HUMAN, FLJ45187, Protein DLN-1. Background: C10orf140, also known as DLN-1, is an 827 amino acid protein that belongs to the DACH/dachshund family. C10orf140 contains a poly-Alanine region that is highly polymorphic.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1404601 | 03/06/2024 | Copyright antibodies-online. All rights reserved.

million base pairs, makes up approximately 4.5 % of total DNA in cells and encodes genes. Several protein-coding genes, including those that encode for chemokines, c excision repair proteins, early growth response factors (Egrs) and fibroblast growth (FGFRs), are located on chromosome 10. Defects in some of the genes that map to chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss: Usher syndrome, nonsyndromatic deafness, Wolman?s syndrome, Cowden syndrom endocrine neoplasia type 2 and porphyria. Gene ID: 387640 Application Details WB 1:300-5000 IHC-P 1:200-400 IHC-P 1:200-400 Restrictions: For Research Use only Handling Iuquid Concentration: 1 µg/µL. Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proc 50 % Glycerol. Preservative: ProClin		
genes. Several protein-coding genes, including those that encode for chemokines, o excision repair proteins, early growth response factors (Egrs) and fibroblast growth (FGFRs), are located on chromosome 10. Defects in some of the genes that map to chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss Usher syndrome, nonsyndromatic deafness, Wolman?s syndrome, Cowden syndrom ene ID: 387640 Application Details Application Notes: WB 1:300-5000 IHC-P 1:200-400 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proson 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.		The gene encoding C10orf140 maps to human chromosome 10, which spans nearly 135
excision repair proteins, early growth response factors (Egrs) and fibroblast growth (FGFRs), are located on chromosome 10. Defects in some of the genes that map to chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss Usher syndrome, nonsyndromatic deafness, Wolman?s syndrome, Cowden syndror endocrine neoplasia type 2 and porphyria. Gene ID: 387640 Application Details WB 1:300-5000 IHC-P 1:200-400 IHC-P 1:200-400 Restrictions: For Research Use only Handling Iuquid Concentration: 1 µg/µL. Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proson % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.		million base pairs, makes up approximately 4.5 % of total DNA in cells and encodes nearly 1,20
(FGFRs), are located on chromosome 10. Defects in some of the genes that map to chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss Usher syndrome, nonsyndromatic deafness, Wolman?s syndrome, Cowden syndrom endocrine neoplasia type 2 and porphyria. Gene ID: 387640 Application Details 387640 Application Notes: WB 1:300-5000 IHC-P 1:200-400 Restrictions: For Research Use only Handling Format: Liquid Concentration: Queous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proson 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.		genes. Several protein-coding genes, including those that encode for chemokines, cadherins,
chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss Usher syndrome, nonsyndromatic deafness, Wolman?s syndrome, Cowden syndromendocrine neoplasia type 2 and porphyria. Gene ID: 387640 Application Details Application Notes: WB 1:300-5000 IHC-P 1:200-400 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proc 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.		excision repair proteins, early growth response factors (Egrs) and fibroblast growth receptors
Usher syndrome, nonsyndromatic deafness, Wolman?s syndrome, Cowden syndrom, cowden syndrome, cowden syndrom, cowden syndrome, co		(FGFRs), are located on chromosome 10. Defects in some of the genes that map to
endocrine neoplasia type 2 and porphyria. Gene ID: 387640 Application Details Application Notes: WB 1:300-5000 IHC-P 1:200-400 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pro 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.		chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss syndrome,
Gene ID: 387640 Application Details International and the second sec		Usher syndrome, nonsyndromatic deafness, Wolman?s syndrome, Cowden syndrome, multipl
Application Details Application Notes: WB 1:300-5000 HC-P 1:200-400 Restrictions: For Research Use only Handling Implied Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pro- 50 % Glycerol. Preservative: ProClln Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.		endocrine neoplasia type 2 and porphyria.
Application Notes: WB 1:300-5000 IHC-P 1:200-400 Restrictions: For Research Use only Handling Iuquid Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pro 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.	ene ID:	387640
IHC-P 1:200-400 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pro 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.	pplication Details	
Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pro- 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.	pplication Notes:	WB 1:300-5000
Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pro 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.		IHC-P 1:200-400
Format:LiquidConcentration:1 μg/μLBuffer:Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pro 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.	estrictions:	For Research Use only
Concentration:1 μg/μLBuffer:Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pro 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.	landling	
Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pro 50 % Glycerol. 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.	ormat:	Liquid
50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.	oncentration:	1 µg/µL
Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.	uffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 an 50 % Glycerol.
handled by trained staff only.	reservative:	ProClin
Storage: -20 °C	recaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
	torage:	-20 °C
Storage Comment: Store at -20°C for 12 months.	torage Comment:	Store at -20°C for 12 months.
Expiry Date: 12 months	xpiry Date:	12 months