



[Go to Product page](#)

Datasheet for ABIN1405029
anti-CLNS1A antibody (AbBy Fluor® 350)

Overview

Quantity:	100 µL
Target:	CLNS1A
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CLNS1A antibody is conjugated to AbBy Fluor® 350
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CLNS1A/pICln
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	CLNS1A
Alternative Name:	CLNS1A (CLNS1A Products)
Background:	Synonyms: nucleotide sensitive 1A, Chloride channel, Chloride channel nucleotide sensitive 1A, Chloride channel regulatory protein, Chloride conductance regulatory protein ICln, Chloride ion current inducer protein, CICI, CLNS 1A, Clns1a, CLNS1B, ICln, ICln, ICLN_HUMAN, Methylosome subunit pICln, Reticulocyte pICln

Target Details

Background: The formation of the spliceosome includes the assembly of Sm proteins in an ordered manner onto snRNAs. This process is mediated by the survival of motor neuron (SMN) protein, and is enhanced by modification of specific arginine residues in the Sm proteins to symmetrical dimethylarginines (sDMAs). sDMA modification of Sm proteins is catalyzed by the methylosome, a complex comprised of the type II methyltransferase PRMT5 (also designated Jak-binding protein 1, JBP1), pICln, and two novel factors. PRMT5 binds the Sm proteins via their arginine- and glycine-rich (RG) domains, while pICln binds the Sm domains. pICln also acts as an inhibitor of SnRNP assembly by preventing specific interactions between Sm proteins required for the formation of the Sm core. pICln is a highly conserved, ubiquitously expressed protein that localizes primarily to the cytoplasm, and may play a role as a swelling-activated anion channel or a channel regulator in addition to its function in the methylosome. The gene encoding human pICln maps to chromosome 11q14.1.

Gene ID: 1207

Pathways: [Ribonucleoprotein Complex Subunit Organization](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months