# antibodies .- online.com





## anti-CLNS1A antibody (AbBy Fluor® 555)



Go to Product page

| ( ) | 11/0   | K\ /   | iew         |
|-----|--------|--------|-------------|
|     | $\cup$ | 'I V/I | $I \cap VV$ |
|     |        |        |             |

| Quantity:    | 100 μL                                                                          |
|--------------|---------------------------------------------------------------------------------|
| Target:      | CLNS1A                                                                          |
| Reactivity:  | Human, Mouse, Rat                                                               |
| Host:        | Rabbit                                                                          |
| Clonality:   | Polyclonal                                                                      |
| Conjugate:   | This CLNS1A antibody is conjugated to AbBy Fluor® 555                           |
| Application: | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

## **Product Details**

| Immunogen:        | KLH conjugated synthetic peptide derived from human CLNS1A/pICln |  |
|-------------------|------------------------------------------------------------------|--|
| Isotype:          | IgG                                                              |  |
| Cross-Reactivity: | Human, Mouse, Rat                                                |  |
| Purification:     | Purified by Protein A.                                           |  |

## Target Details

| Target:                                                                                                                                                                                                                                                     | CLNS1A                                                                                                           |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------|
| Alternative Name:                                                                                                                                                                                                                                           | CLNS1A (CLNS1A Products)                                                                                         |
| Background: Synonyms: nucleotide sensitive 1A, Chloride channel, Chloride channel nucleotide Chloride channel regulatory protein, Chloride conductance regulatory protein IC current inducer protein, CICI, CLNS 1A, Clns1a, CLNS1B, ICIn, ICIn, ICLN_HUMAN |                                                                                                                  |
|                                                                                                                                                                                                                                                             | current inducer protein, CICI, CLNS 1A, Clns1a, CLNS1B, ICIn, ICIn, ICLN_HUMAN subunit pICIn, Reticulocyte pICIn |

Background: The formation of the spliceosome includes the assembly of Sm proteins in an ordered manner onto snRNAs. This process is mediated by the survival of motor neuron (SMN) protein, and is enhanced by modification of specific arginine residues in the Sm proteins to symmetrical dimethylarginines (sDMAs). sDMA modification of Sm proteins is catalyzed by the methylosome, a complex comprised of the type II methyltransferase PRMT5 (also designated Jak-binding protein 1, JBP1), plCln, and two novel factors. PRMT5 binds the Sm proteins via their arginine- and glycine-rich (RG) domains, while plCln binds the Sm domains. plCln also acts as an inhibitor of SnRNP assembly by preventing specific interactions between Sm proteins required for the formation of the Sm core. plCln is a highly conserved, ubiquitously expressed protein that localizes primarily to the cytoplasm, and may play a role as a swelling-activated anion channel or a channel regulator in addition to its function in the methylosome. The gene encoding human plCln maps to chromosome 11q14.1.

Gene ID:

1207

Pathways:

Ribonucleoprotein Complex Subunit Organization

## **Application Details**

Application Notes:

IF(IHC-P) 1:50-200

Restrictions:

For Research Use only

#### Handling

| Format:            | Liquid                                                                                                             |
|--------------------|--------------------------------------------------------------------------------------------------------------------|
| Concentration:     | 1 μg/μL                                                                                                            |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin                                                                                                            |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C                                                                                                             |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months                                                                                                          |