antibodies

## Datasheet for ABIN1405038 anti-TFCP2 antibody (AbBy Fluor® 647)



Overview	
Quantity:	100 µL
Target:	TFCP2
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TFCP2 antibody is conjugated to AbBy Fluor® 647
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human TFCP2C
lsotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.
Target Details	
Target:	TFCP2
Alternative Name:	TFCP2C (TFCP2 Products)
Background:	Synonyms: Alpha globin transcription factor CP2, CP2, LBP 1C, LBP1C, LSF, SAA3 enhancer factor, SEF, TFCP2, TFCP2C, Transcription factor CP2, Transcription factor LSF, TFCP2_HUMAN.
	Background: CP2 family gene products are reported to be involved in a-globin gene expression.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1405038 | 06/07/2024 | Copyright antibodies-online. All rights reserved.

	Mouse CP2c (also known as a-CP2/CP2/LSF/UBP-1) is a homologue of human LBP-1c with
	96 % amino acid sequence identity. CP2c was initially identified as an activator of the mouse a-
	globin gene, which binds to the consensus DNA-binding sequence CNRG-N6-CNR(G/C) within a
	promoter element overlapping the CCAAT box. The transcriptional activity of CP2c was
	increased during induced differentiation of MEL cells and associated globin gene transcription.
	CP2c is also likely to be involved in regulation of non-globin erythroid-specific genes. It was
	reported that congenital erythropoietic porphyria was caused by the mutations in GATA-1 and
	CP2c binding sites within the promoter of the uroporphyrinogen III synthase gene, the fourth enzyme in the heme biosynthetic pathway.
Gene ID:	7024
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months