# antibodies -online.com





Datasheet for ABIN1405039

# anti-TFCP2 antibody (Biotin)



$\sim$			
	N/P	r\/I	i⊢₩

Quantity:	100 μL	
Target:	TFCP2	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TFCP2 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human TFCP2C	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	

# Target Details

Target:	TFCP2
Alternative Name:	TFCP2C (TFCP2 Products)
Background:	Synonyms: Alpha globin transcription factor CP2, CP2, LBP 1C, LBP1C, LSF, SAA3 enhancer factor, SEF, TFCP2, TFCP2C, Transcription factor CP2, Transcription factor LSF,
	TFCP2_HUMAN.  Background: CP2 family gene products are reported to be involved in a-globin gene expression.

#### **Target Details**

Mouse CP2c (also known as a-CP2/CP2/LSF/UBP-1) is a homologue of human LBP-1c with 96 % amino acid sequence identity. CP2c was initially identified as an activator of the mouse a-globin gene, which binds to the consensus DNA-binding sequence CNRG-N6-CNR(G/C) within a promoter element overlapping the CCAAT box. The transcriptional activity of CP2c was increased during induced differentiation of MEL cells and associated globin gene transcription. CP2c is also likely to be involved in regulation of non-globin erythroid-specific genes. It was reported that congenital erythropoietic porphyria was caused by the mutations in GATA-1 and CP2c binding sites within the promoter of the uroporphyrinogen III synthase gene, the fourth enzyme in the heme biosynthetic pathway.

Gene ID:

7024

## **Application Details**

Application Notes: WB 1:300-5000 IHC-P 1:200-400

Restrictions: For Research Use only

### Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C for 12 months.	
Expiry Date:	12 months	