antibodies -online.com





anti-Kcng2 antibody (Alexa Fluor 350)



()	1 /	0	rv	/ 1 /	71	Α.
	1//	-	1 \/	16		1/1/
\sim	v	\sim	1 V	١,	_	v v

Quantity:	100 μL	
Target:	Kcng2	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Kcng2 antibody is conjugated to Alexa Fluor 350	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human KCNG2
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	Kcng2
Alternative Name:	KCNG2 (Kcng2 Products)
Background:	Synonyms: Cardiac potassium channel subunit, KCNG2, KCNG2_HUMAN, KV6.2, Potassium voltage-gated channel subfamily G member 2, Voltage-gated potassium channel subunit Kv6.2. Background: Neuronal and cardiac cells are excited by voltage-gated ion channels. Voltage-
	gated K+ channels in the plasma membrane control the repolarization and the frequency of

action potentials in neurons, muscles and other excitable cells. Mutations interfering with potassium ion channels are known to cause a variety of disorders. KCNG2 (potassium voltage-gated channel subfamily G member 2) is also known as voltage-gated potassium channel subunit KV6.2, cardiac potassium channel subunit or KCNF2 and is a 466 amino acid protein. KCNG2 is a multi-pass membrane protein abundantly expressed in heart, liver, skeletal muscle, kidney and pancreas, and detected at lower concentrations in brain, lung and placenta. KCNG2 is an electrically silent subunit that forms heterodimers with KV2.1, creating a unique functional K+ channel. KCNG2-KV2.1 heterodimers are known to be inhibited by tetraethylammonium and propafenone. KCNG2 is thought to downregulate potassium channel currents because KCNG2-KV2.1 heterodimers generate smaller currents than KV2.1 homodimers

Gene ID:

26251

UniProt:

Q9UJ96

Application Details

Application Notes:	IF(IHC-P) 1:50-200
, ippiiodition i totooi	() = 200

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months