antibodies .- online.com





anti-CLEC11A antibody (AbBy Fluor® 647)



()	ve	K\ /		A .
	\cup	1 V/	Щ.	V۷

Overview		
Quantity:	100 μL	
Target:	CLEC11A	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CLEC11A antibody is conjugated to AbBy Fluor® 647	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
	KLH conjugated synthetic peptide derived from human Pleckstrin/p47	
Immunogen:	KLH conjugated synthetic peptide derived from human Pleckstrin/p47	
Immunogen: Isotype:	KLH conjugated synthetic peptide derived from human Pleckstrin/p47	
Isotype:	IgG	
Isotype: Cross-Reactivity:	IgG Human, Mouse, Rat	
Isotype: Cross-Reactivity: Purification:	IgG Human, Mouse, Rat	
Isotype: Cross-Reactivity: Purification: Target Details	IgG Human, Mouse, Rat Purified by Protein A.	

Background: Platelet P47, also known as NSFL1C, UBX1, UBXD10 or UBXN2C, is a 370 amino

acid protein that localizes to both the nucleus and the golgi apparatus (specifically to golgi

Target Details

stacks) and contains one SEP domain and one UBX domain. Functioning as part of a ternary complex with VCP (a protein involved in the heterotypic fusion of transport vesicles with their target membranes) and Syntaxin 5, p47 interacts with and reduces the ATPase activity of VCP and is required for the fragmentation of golgi stacks during mitosis and for subsequent reassembly of golgi stacks after mitosis. p47 is subject to phosphorylation during mitosis, which inhibits p47-golgi interaction and is, therefore, required for proper golgi stack formation and cisternal regrowth. Human p47 shares 89 % sequence identity with its mouse counterpart, suggesting a conserved role between species. Multiple isoforms of p47 exist due to alternative splicing events.

Gene ID:

5341

Application Details

Application Notes:	IF(IHC-P) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	