## antibodies .- online.com





## anti-GPLD1 antibody (Biotin)



Go to Product page

( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL	
Target:	GPLD1	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This GPLD1 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	
Product Details		
1 Toddet Details		
Immunogen:	KLH conjugated synthetic peptide derived from human GPLD1/GPI-PLD	
	KLH conjugated synthetic peptide derived from human GPLD1/GPI-PLD	
Immunogen:		
Immunogen: Isotype:	IgG	
Immunogen: Isotype: Cross-Reactivity:	IgG Human, Mouse, Rat	
Immunogen: Isotype: Cross-Reactivity: Purification:	IgG Human, Mouse, Rat	
Immunogen: Isotype: Cross-Reactivity: Purification: Target Details	lgG Human, Mouse, Rat Purified by Protein A.	

phospholipase D, PI-G PLD, PIGPLD, PIGPLD1.

D, GPI-PLD, GPI-specic phospholipase D, GPIPLD, GPIPLDM, GPLD1, PHLD\_HUMAN,

Phosphatidylinositol glycan specic phospholipase D, Phosphatidylinositol-glycan-specic

Background: Phosphatidylinositol-glycan-specific phospholipase D (GPI-PLD) is a high-density lipoprotein-associated protein found on chromosome 6p22 that specifically hydrolyzes the inositol phosphate linkage in proteins anchored by phosphatidylinositol-glycans (PI-Gs). GPI-PLD is found in serum, liver, cerebrospinal fluid and in milk. The majority of plasma GPI-PLD appears to be specifically associated with a small, discrete and minor fraction of lipoproteins containing apoA-I and apoA-IV. Serum GPI-PLD activity is reduced over 75 % in systemic inflammatory response syndrome and the downregulation of GPI-PLD could play an important role in the control of proinflammatory responses.

Gene ID:

2822

Pathways:

Inositol Metabolic Process, Regulation of Carbohydrate Metabolic Process

## **Application Details**

Application Notes:

WB 1:300-5000

IHC-P 1:200-400

12 months

Restrictions:

For Research Use only

## Handling

**Expiry Date:** 

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C for 12 months.	