Datasheet for ABIN1405424

anti-PAPPA2 antibody (FITC)

Go to Product page

Overview

Quantity:	100 µL
Target:	PAPPA2
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PAPPA2 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PAPP A2
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	PAPPA2
Alternative Name:	PAPP A2 (PAPPA2 Products)
Background:	Synonyms: PAPPA 2, PAPP E, PAPP-A2, PAPP-E, PAPP2_HUMAN, PAPPA 2, PAPPA2, Pappalysin 2, Pappalysin-2, PLAC3, Pregnancy associated plasma protein A2, Pregnancy associated plasma protein E1, Pregnancy-associated plasma protein A2, Pregnancy-associated plasma protein E1.

Copyright antibodies-online. All rights reserved.

Target Details

	Background: Pregnancy-associated plasma protein-A (Pappalysin-1 or PAPP-A), also known as Insulin-like growth factor-dependent IGF-binding protein 4 (IGFBP4) protease, is a member of the peptidase M43B family of proteins. PAPP-A, a metalloproteinase, cleaves Insulin-like growth factor binding proteins IGFBP4 and IGFBP5, releasing bound IGF. PAPP-A is primarily expressed in septa and anchoring villi in placenta and is also expressed in pregnancy serum. Levels of PAPP-A increase throughout pregnancy. Lower levels of expression can be detected in kidney, prostate, breast, ovary and endometrial tissues. PAPP-A is a secreted protein that can form homodimers, in pregnancy serum PAPP-A may also form a heterotetramer with PRG-2.
Gene ID:	60676

Application Details

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months

Copyright antibodies-online. All rights reserved.