

[Go to Product page](#)

Datasheet for ABIN1406757

**anti-GLDN antibody (AA 365-460) (Cy3)**

## Overview

Quantity:	100 µL
Target:	GLDN
Binding Specificity:	AA 365-460
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GLDN antibody is conjugated to Cy3
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Gliomedin/COLM
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

## Target Details

Target:	GLDN
Alternative Name:	Gliomedin ( <a href="#">GLDN Products</a> )

## Target Details

Background:	<p>Synonyms: CANCER RELATED GENE LIVER 2, Cancer related gene-Liver 2, CLOM, collomin, Colm gene name, COLM, Colmedin, CRG L2, FLJ23917, Gldn gene name, UNC 112, GLDN_HUMAN.</p> <p>Background: Gliomedin is a 551 amino acid protein encoded by the human gene GLDN. Gliomedin is thought to play a role in the formation of the nodes of Ranvier along myelinated axons. Accumulation of Na<sup>+</sup> channels at the nodes of Ranvier is a prerequisite for saltatory conduction. In peripheral nerves, clustering of these channels along the axolemma is regulated by myelinating Schwann cells through an unknown mechanism. Gliomedin is a glial ligand for Neurofascin and NrCAM, two axonal immunoglobulin cell adhesion molecules that are associated with Na<sup>+</sup> channels at the nodes of Ranvier. Gliomedin is expressed by myelinating Schwann cells and accumulates at the edges of each myelin segment during development, where it aligns with the forming nodes. Gliomedin is a single-pass type II membrane protein localized to the nodes of Ranvier and is specifically expressed in spinal cord, brain, placenta and sciatic nerve. It is more abundant in peripheral than central nervous system.</p>
-------------	--

## Application Details

Application Notes:	<p>IF(IHC-P) 1:50-200</p> <p>IF(IHC-F) 1:50-200</p> <p>IF(ICC) 1:50-200</p>
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months