# antibodies - online.com







# anti-Astrotactin 2 antibody (AA 21-120) (HRP)



( )	1/0	r\ / I	014	
( )	ve	I V I	-v	V

Quantity:	100 μL	
Target:	Astrotactin 2 (ASTN2)	
Binding Specificity:	AA 21-120	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Astrotactin 2 antibody is conjugated to HRP	
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human ASTN2
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow
Purification:	Purified by Protein A.

### **Target Details**

Target:	Astrotactin 2 (ASTN2)
Alternative Name:	ASTN2 (ASTN2 Products)
Background: Synonyms: Astn2, Astn-2, Astn 2, ASTN2_HUMAN, Astrotactin 2, Astrotactin2, Astrotactin-	

bA67K19.1, KIAA0634.

For Research Use only

Background: Neuronal migration of the central nervous system is a specialized form of cell motility that takes place in the laminar structure of the cortical regions of brain. Astrotactin is a neuronal cell surface protein expressed on postmitotic neuronal precursors in the cerebellum, hippocampus, cerebrum, and olfactory bulb. Astrotactin 2 is a paralog of Astrotactin. Astrotactin mediates neuron-astroglial interactions and is also implicated in synaptic development as well as many other neuronal activities. Astrotactin has three epidermal growth factor repeat domains and two fibronectin type III repeat domains. The human Astrotactin gene shows extensive homology to the mouse Astrotactin gene. Mutations in the Astrotactin gene are linked to neuronal migration defects in both species.

#### **Application Details**

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500

## Handling

Restrictions:

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months