

[Go to Product page](#)

Datasheet for ABIN1407184

**anti-Neurexin 2 (NRXN2) (AA 971-1070) antibody (Cy5)**

## Overview

Quantity:	100 µL
Target:	Neurexin 2 (NRXN2)
Binding Specificity:	AA 971-1070
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	Cy5
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NRXN2
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human, Mouse, Dog, Cow, Sheep, Pig, Horse
Purification:	Purified by Protein A.

## Target Details

Target:	Neurexin 2 (NRXN2)
Alternative Name:	Neurexin 2 ( <a href="#">NRXN2 Products</a> )

## Target Details

Background:	<p>Synonyms: Neurexin 2, Neurexin-2, Neurexin2, Neurexin II, Neurexin II alpha, NRXN 2, NRXN-2, NRXN2, NRX2A_HUMAN.</p> <p>Background: Neurexins comprise a family of neuronal cell surface proteins, which include neurexin I (NRXN1), neurexin II (NRXN2), neurexin III (NRXN3) and Caspr (neurexin IV). Neurexins I-III are expressed as a and b isoforms. The a isoforms are made of three cassettes, which contain two LNS (Laminin A, Neurexins, Sex hormone-binding)-domains separated by EGF domains, followed by a transmembrane region and a 55 amino acid cytoplasmic C-terminal. The a isoforms bind to neurexophilins at the second LNS site and to the excitatory neurotoxin <math>\alpha</math>-latrotoxin. The b isoforms have only one LNS-domain, bind to neuroligins, and play a role in the formation and remodeling of synapses. Caspr (for Contactin-Associated Protein 1, also designated Paranodin in mouse), contains an extracellular domain similar to the other three neurexins, and binds to the surface glycoprotein Contactin. Caspr and the closely related Caspr2, a mammalian homolog of Drosophila Neurexin IV (Nrx-IV), demarcate distinct subdomains in myelinated axons. Specifically, Caspr exists at the paranodal junctions, while Caspr2 colocalizes with Shaker-like K<sup>+</sup> channels in the juxtaparanodal region. Caspr may play a role in the communication of glial cells and neurons during development.</p>
-------------	--

UniProt:	<a href="#">Q9P2S2</a>
----------	------------------------

Pathways:	<a href="#">Synaptic Membrane</a>
-----------	-----------------------------------

## Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

---

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months