.-online.com antibodies

## Datasheet for ABIN1408282 anti-HAS2 antibody (Cy5)

Background:



Overview	
Quantity:	100 μL
Target:	HAS2
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HAS2 antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human HAS2/Hyaluronan synthase 2
lsotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.
Target Details	
Target:	HAS2
Alternative Name:	Has2 (HAS2 Products)

2, Hyaluronic acid synthase 2.

Background: HAS1, HAS2 and HAS3 are HA Synthase proteins that synthesize HA (Hyaluronan

Synonyms: HA synthase 2, has2, HAS2\_HUMAN, Hyaluronan synthase 2, Hyaluronate synthase

or hyaluronic acid). The extracellular matrix in most vertebrates express HA, which is a high

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1408282 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

	molecular weight linear polysaccharide composed of alternating glucuronic acid and N-
	acetylglucosamine residues linked by i²-1,3 and i²-1,4 glycosidic bonds. The three HAS genes
	show distinct patterns of expression during development and their protein products play
	significantly different roles in the formation of the HA matrix. Both HAS1 and HAS2 synthesise
	high molecular-weight HA, whereas HAS3 produces lower molecular weight HA. The expression
	of the three HAS isoforms is more prominent in growing cells than in resting cells and is
	differentially regulated by various stimuli suggesting distinct functional roles of the three
	proteins. HAS2 mRNA shows predominant expression in chondrocytes and cartilage. The
	human HAS2 gene maps to chromosome 8q24.12.
Pathways:	Glycosaminoglycan Metabolic Process
Application Details	
Application Notes:	IF(IHC-P): (1:50-200)
	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.