.-online.com antibodies

## Datasheet for ABIN1408389 anti-SRD5A1 antibody (Cy3)



Overview 100 µL Quantity: Target: SRD5A1 Reactivity: Human, Rat, Mouse Host: Rabbit Clonality: Polyclonal Conjugate: This SRD5A1 antibody is conjugated to Cy3 Application: Western Blotting (WB) **Product Details** Immunogen: KLH conjugated synthetic peptide derived from human SRD5A1 Isotype: IgG Cross-Reactivity: Cow, Human, Sheep Predicted Reactivity: Mouse,Rat,Dog,Horse,Chicken Purification: Purified by Protein A.

## Target Details

Target:	SRD5A1
Alternative Name:	Srd5a1 (SRD5A1 Products)
Background:	Synonyms: 3-oxo-5-alpha-steroid 4-dehydrogenase 1, SR type 1, Steroid 5-alpha-reductase 1, S5AR 1, S5A1_HUMAN.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1408389 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

	Background: Steroid 5a-Reductase is an important enzyme in androgen physiology because it
	catalyzes the conversion of testosterone into the more potent 5a-dihydro-testosterone, which
	mediates androgen effects on target tissues. The enzyme exists as two isoforms: type 1, which
	is expressed mainly in the skin, and type 2, which is expressed mainly in the prostate. In
	cultured human skin cells, 5a-Reductase 1 shows heterogeneity of protein, and has different
	levels of transcriptional and translational expression. 5a-Reductase 1 is expressed in all
	portions of the hair follicle, whereas 5a-Reductase 2 is expressed only in mesenchymal
	portions. In addition, 5a-Reductase 1 is mainly expressed in human breast carcinoma and may
	play a role in the in situ production and actions of the potent androgen 5a-dihydrotestosterone,
	including inhibition of cancer cell proliferation in hormone-dependent human breast carcinoma.
	The 5a-Reductase-3a-hydroxysteroid dehydrogenase complex is present in the human brain,
	suggesting that the complex may be involved in the synthesis of neuroactive steroids or the
	catabolism of neurotoxic steroids.
Pathways:	Metabolism of Steroid Hormones and Vitamin D, Steroid Hormone Biosynthesis, Response to
	Growth Hormone Stimulus, C21-Steroid Hormone Metabolic Process

## Application Details

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN1408389 | 03/07/2024 | Copyright antibodies-online. All rights reserved.