

[Go to Product page](#)

Datasheet for ABIN1408521  
**anti-SORCS1 antibody (AA 561-660) (Cy3)**

## Overview

Quantity:	100 µL
Target:	SORCS1
Binding Specificity:	AA 561-660
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SORCS1 antibody is conjugated to Cy3
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SORCS1
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

## Target Details

Target:	SORCS1
Alternative Name:	SORCS1 ( <a href="#">SORCS1 Products</a> )
Background:	Synonyms: SorCS 1, SORCS, SORCS-1, SORCS receptor 1, Sortilin related VPS10 domain

## Target Details

containing receptor 1, VPS10 domain containing receptor SorCS1, VPS10 domain receptor protein SORCS 1, VPS10 domain receptor protein SORCS, VPS10 domain receptor protein SORCS1, VPS10 domain receptor SorCS, FLJ41758, FLJ43475, FLJ44957, hSorCS, OTTHUMP00000058855, SORC1\_HUMAN.

Background: There are three sorCS genes that have diverse, partially overlapping functions in the central nervous system. In the developing and mature central nervous system, the homologous SorCS1 and SorCS2 genes and the SorCS3 gene are expressed in a combinatorial, non-overlapping pattern. SorCS proteins show homology to the mosaic receptor SorLA and the neurotensin receptor sortilin, based on a common VPS10 domain, which is the hallmark of the SorCS receptor family. SorCS1 is a type 1 receptor containing a VPS10P domain and a leucine-rich domain. Alternative splicing of human SorCS1 results in four isoforms with different cytoplasmic tails and differential expression in tissues. Human SorCS1 is detected in fetal and infant brain and in fetal retina. Alternative splicing of murine SorCS1 also results in four isoforms. Murine isoform 1 is highly expressed in brain and at lower levels in heart, liver and kidney. It is detected in newborn mouse brain and in adult olfactory bulb and cerebral cortex. Murine isoform 2 is highly expressed in liver and at lower levels in heart, brain, kidney and testis.

## Application Details

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Handling

---

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months