

Datasheet for ABIN1408638

anti-ICA1 antibody (AA 1-100) (Cy7)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	ICA1
Binding Specificity:	AA 1-100
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ICA1 antibody is conjugated to Cy7
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ICA1
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Cow, Pig, Horse, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	ICA1
Alternative Name:	ICA1 (ICA1 Products)
Background:	Synonyms: 69 kDa islet cell autoantigen, Diabetes mellitus type I autoantigen, ICA 1, ICA1,

Target Details

ICA69, ICA69_HUMAN, ICAp69, Islet cell autoantigen 1 69kD, Islet cell autoantigen 1 69 kDa, Islet cell autoantigen 1, Islet cell autoantigen 1 isoform, Islet cell autoantigen p69, OTTHUMP00000200933, OTTHUMP00000200934, OTTHUMP00000200941, OTTHUMP00000200993, p69.

Background: Carbonic anhydrases (CAs), also designated carbonate dehydratases or carbonate hydrolyases, form a large family of genes that encode zinc metalloenzymes of great physiologic importance. As catalysts of the reversible hydration of carbon dioxide, these enzymes participate in a variety of biologic processes, including respiration, acid-base balance, bone resorption and calcification as well as the formation of aqueous humor, cerebrospinal fluid, saliva and gastric acid. Genes in the ?carbonic anhydrase family encode either active carbonic anhydrase isozymes or _catalytic?(devoid of CO₂ hydration activity) carbonic anhydrase-related proteins. Human CA I (CA1) is encoded by the CA1 gene, which maps to a region on chromosome 8 that harbors a cluster of CA genes. CA I localizes to the cytoplasm and research indicates that a severe deficiency of CA I does not result in any obvious hematological or renal consequences.

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months