Datasheet for ABIN1408690 anti-RIMBP2 antibody (AA 951-1052) (Cy5)

-online.com antibodies



\sim		
$() \setminus I \subset$	1/1/1	$\Box \wedge A$
Ove	71 V I	C V V

Quantity:	100 µL
Target:	RIMBP2
Binding Specificity:	AA 951-1052
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RIMBP2 antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human RIMBP2
Isotype:	lgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	RIMBP2
Alternative Name:	RIMBP2 (RIMBP2 Products)
Background:	Synonyms: RIM-BP2, RIMB2_HUMAN, RIMBP2, RIMS-binding protein 2.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1408690 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

release by forming links between synaptic-vesicle fusion apparatuses and calcium cha Specifically, RIM-BP2 (RIMS binding protein 2), also known as RBP2, is a 1,052 amino. protein that links L-type Ca++ CP Alpha1D, N-type Ca++ CP Alpha1B, Rim1 and Rim2 or synaptic transmission. RIM-BP2 contains three fibronectin type-III domains and three domains, which are used to mediate binding to a proline-rich motifs. Existing as three alternatively spliced isoforms, RIM-BP2 is encoded by a gene that maps to human chromosome 12q24.33 and mouse chromosome 5 G1.3. Gene ID: 23504 Application Details Application Notes: IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 Buffer: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered soluti		
Specifically, RIM-BP2 (RIMS binding protein 2), also known as RBP2, is a 1,052 amino. protein that links L-type Ca++ CP Alpha1D, N-type Ca++ CP Alpha1B, Rim1 and Rim2 c synaptic transmission. RIM-BP2 contains three fibronectin type-III domains and three domains, which are used to mediate binding to a proline-rich motifs. Existing as three alternatively spliced isoforms, RIM-BP2 is encoded by a gene that maps to human chromosome 12q24.33 and mouse chromosome 5 G1.3. Gene ID: 23504 Application Details Application Notes: IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(IHC-F) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 IF(UCC) 1:50-200 IF(UCC) 1:50-200 IF(UCC) 1:50-200 IF(UCC) 1:50-200 Restrictions: For Research Use only Handling Eormat: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Prool 50 % Glycerol. Preservative: ProClin Preservative: ProClin		Background: RIMS-binding proteins (RIM-BPs) serve as adaptors during vesicle fusion and
protein that links L-type Ca++ CP Alpha1D, N-type Ca++ CP Alpha1B, Rim1 and Rim2 c synaptic transmission. RIM-BP2 contains three fibronectin type-III domains and three domains, which are used to mediate binding to a proline-rich motifs. Existing as three alternatively spliced isoforms, RIM-BP2 is encoded by a gene that maps to human chromosome 12q24.33 and mouse chromosome 5 G1.3. Gene ID: 23504 Application Details Application Notes: IF(IHC-P) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Eormat: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % ProcI 50 % Glycerol. Proclin Preservative: ProClin		release by forming links between synaptic-vesicle fusion apparatuses and calcium channels.
synaptic transmission. RIM-BP2 contains three fibronectin type-III domains and three domains, which are used to mediate binding to a proline-rich motifs. Existing as three alternatively spliced isoforms, RIM-BP2 is encoded by a gene that maps to human chromosome 12q24.33 and mouse chromosome 5 G1.3. Gene ID: 23504 Application Details IF(IHC-P) 1:50-200 IF(ICC) IF(ICC) 1:50-200 IF(ICC) IF(ICC) 1:50-200 IF(ICC) IF(ICC) 1:50-200 IF(ICC) IF(Specifically, RIM-BP2 (RIMS binding protein 2), also known as RBP2, is a 1,052 amino acid
domains, which are used to mediate binding to a proline-rich motifs. Existing as three alternatively spliced isoforms, RIM-BP2 is encoded by a gene that maps to human chromosome 12q24.33 and mouse chromosome 5 G1.3. Gene ID: 23504 Application Details IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(IHC-F) 1:50-200 IF(IHC-F) 1:50-200 IF(IHC-F) 1:50-200 Restrictions: For Research Use only Handling If(IHC-F) 1:50-200 Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procided S0 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which shandled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		protein that links L-type Ca++ CP Alpha1D, N-type Ca++ CP Alpha1B, Rim1 and Rim2 during
alternatively spliced isoforms, RIM-BP2 is encoded by a gene that maps to human chromosome 12q24.33 and mouse chromosome 5 G1.3. Gene ID: 23504 Application Details IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(IHC-F) 1:50-200 IF(IHC-F) 1:50-200 IF(IHC-F) 1:50-200 Restrictions: For Research Use only Handling If(IHC-F) 1:50-200 Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procells 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which shandled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		synaptic transmission. RIM-BP2 contains three fibronectin type-III domains and three SH3
chromosome 12q24.33 and mouse chromosome 5 G1.3. Gene ID: 23504 Application Details Application Notes: IF(IHC-P) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proof 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which a handled by trained staff only. Storage -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		domains, which are used to mediate binding to a proline-rich motifs. Existing as three
Gene ID: 23504 Application Details Application Notes: IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Processon 50 % Glycerol. Preservative: Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which shandled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		alternatively spliced isoforms, RIM-BP2 is encoded by a gene that maps to human
Application Details Application Notes: IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Image: Concentration: Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procells 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which e handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		chromosome 12q24.33 and mouse chromosome 5 G1.3.
Application Notes: IF(IHC-P) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Iuiuid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proof 50 % Glycerol. Preservative: Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which shandled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Gene ID:	23504
IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procells 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which is handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Application Details	
IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proctions 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which shandled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Application Notes:	IF(IHC-P) 1:50-200
Restrictions: For Research Use only Handling Liquid Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proof 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which is handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		IF(IHC-F) 1:50-200
HandlingFormat:LiquidConcentration:1 μg/μLBuffer:Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proof 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which s handled by trained staff only.Storage:-20 °CStorage Comment:Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		IF(ICC) 1:50-200
Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclis 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which is handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Restrictions:	For Research Use only
Concentration:1 μg/μLBuffer:Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclis 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which s handled by trained staff only.Storage:-20 °CStorage Comment:Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Handling	
Buffer:Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procli 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which s handled by trained staff only.Storage:-20 °CStorage Comment:Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Format:	Liquid
50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which shandled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Concentration:	1 μg/μL
Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which is handled by trained staff only.Storage:-20 °CStorage Comment:Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which is handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		50 % Glycerol.
handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Preservative:	ProClin
Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Precaution of Use:	
Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	Storage:	handled by trained staff only.
Expiry Date: 12 months		handled by trained staff only. -20 °C

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN1408690 | 03/07/2024 | Copyright antibodies-online. All rights reserved.