# antibodies -online.com







# anti-RIMBP2 antibody (AA 951-1052) (HRP)



( )	11	$\sim$	rv		۱ ۸
	1 \ /	⊢	I \/	╙	1/1

Quantity:	100 μL	
Target:	RIMBP2	
Binding Specificity:	AA 951-1052	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This RIMBP2 antibody is conjugated to HRP	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human RIMBP2
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Horse,Chicken
Purification:	Purified by Protein A.

## **Target Details**

Target:	RIMBP2	
Alternative Name:	RIMBP2 (RIMBP2 Products)	
Background:	Synonyms: RIM-BP2, RIMB2_HUMAN, RIMBP2, RIMS-binding protein 2.	

#### **Target Details**

Background: RIMS-binding proteins (RIM-BPs) serve as adaptors during vesicle fusion and release by forming links between synaptic-vesicle fusion apparatuses and calcium channels. Specifically, RIM-BP2 (RIMS binding protein 2), also known as RBP2, is a 1,052 amino acid protein that links L-type Ca++ CP Alpha1D, N-type Ca++ CP Alpha1B, Rim1 and Rim2 during synaptic transmission. RIM-BP2 contains three fibronectin type-III domains and three SH3 domains, which are used to mediate binding to a proline-rich motifs. Existing as three alternatively spliced isoforms, RIM-BP2 is encoded by a gene that maps to human chromosome 12q24.33 and mouse chromosome 5 G1.3.

Gene ID:

23504

For Research Use only

### **Application Details**

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500

# Handling

Restrictions:

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months