# antibodies - online.com







# anti-RIMS4 antibody (AA 1-100) (Cy7)



Go t	to I	<sup>2</sup> rod	uct	page	٥

( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL
Target:	RIMS4
Binding Specificity:	AA 1-100
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RIMS4 antibody is conjugated to Cy7
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human RIMS4
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Horse,Rabbit
Purification:	Purified by Protein A.

# **Target Details**

Target:	RIMS4	
Alternative Name:	RIMS4 (RIMS4 Products)	
Background:	Synonyms: C20orf190, dJ781B1.3, Rab3-interacting molecule 4, regulating synaptic membra	

exocytosis 4, Regulating synaptic membrane exocytosis protein 4, RIM 4, RIM4 gamma, Rims4,
RIMS4 HUMAN.

Background: Rim4 (Rab 3 interacting molecule 4), also known as Rim4 or regulating synaptic membrane exocytosis protein 4, is a 269 amino acid protein that localizes to the cell junction and regulates synaptic membrane exocytosis. Rab 3, a neural/neuroendocrine-specific member of the Rab family, is involved in Ca2+-regulated exocytosis. Rab 3 functions in an inhibitory capacity by controlling the recruitment of secretory vesicles into a releasable pool at the plasma membrane. Rim (Rab 3 interacting molecule), a putative effector protein for Rab 3 proteins, is thought to regulate neutrotransmitter release through its interaction with Rab 3 and other synaptic proteins.

Gene ID:	140730

Pathways: Synaptic Vesicle Exocytosis

## **Application Details**

Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	

Restrictions: For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months