

## Datasheet for ABIN1408805

# anti-Synaptojanin 2 antibody (AA 401-500) (Cy5.5)



### Overview

Overview		
Quantity:	100 μL	
Target:	Synaptojanin 2 (SYNJ2)	
Binding Specificity:	AA 401-500	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Synaptojanin 2 antibody is conjugated to Cy5.5	
Application:	: Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescen	
	(Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human Synaptojanin 2	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat	
Purification:	Purified by Protein A.	
r armoution.	Purmed by Protein A.	
Target Details	Pullied by Protein A.	
	Synaptojanin 2 (SYNJ2)	
Target Details		

INPP5H, KIAA0348, MGC44422, SJ2, Synaptic inositol 1 4 5 trisphosphate 5 phosphatase 2, Synaptic inositol-1, Synaptojanin-2, Synaptojanin2, SYNJ 2, Synj2, SYNJ2\_HUMAN.

Background: The inositol polyphosphate 5-phosphatases selectively remove the phosphate from the 5-position of various phosphatidylinositols, which generate second messengers in response to extracellular signals. Synaptojanins are characterized by an N-terminal SAC1-like sequence, a central 5-phosphate domain, and a unique C-terminal sequence and have been shown to use phosphatidylinositol 4,5-bisphosphate as a substrate. Synaptojanins exist as two isoforms, synaptojanin 1 and 2, which differ in the C-terminal domain, and each isoform has multiple variants produced by alternative splicing. Synaptojanin 1 is expressed as two major forms: the shorter is found in brain while the longer is expressed in peripheral tissues. Eight splice variants of synaptojanin 2 have been detected, including a brain specific isoform. Synaptojanins are thought to participate in the endocytosis of synaptic vesicles and the regulation of the actin cytoskeleton.

Gene ID: 8871

Pathways: Inositol Metabolic Process

#### **Application Details**

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

	_	
-	II	1:
-	-222	nnc
-	Hand	1111()

Expiry Date:

12 months