

[Go to Product page](#)

Datasheet for ABIN1408840

anti-AACS antibody (AA 182-228) (Cy5)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | AACS |
| Binding Specificity: | AA 182-228 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This AACS antibody is conjugated to Cy5 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human AACS |
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|---|
| Target: | AACS |
| Alternative Name: | AACS/ACSF1 (AACS Products) |
| Background: | Synonyms: Acetoacetyl-CoA synthetase, ACSF1, Acyl-CoA synthetase family member 1, Protein |

Target Details

sur-5 homolog, AACCS_HUMAN.

Background: ACSF1 is a 672 amino acid protein belonging to the ATP-dependent AMP-binding enzyme family. Encoded by a gene that maps to human chromosome 12q24.31, ACSF1 is highly expressed in kidney, heart and brain, and shows similar neural expression as HMGCR (3-hydroxy-3-methylglutaryl-CoA reductase). Existing as three alternatively spliced isoforms, ACSF1 participates in ATP binding, ligase activity, acetoacetate-CoA ligase activity and nucleotide binding. The ACSF1 promoter is a known PPAR α target gene, with the nuclear receptor recruited to the ACSF1 promoter by direct interaction with stimulating protein-1 (Sp1). ACSF1 activates acetoacetate and is highly regulated by modulators that affect HMGCR and cholesterol biosynthesis.

Gene ID: 65985

Pathways: [Positive Regulation of Peptide Hormone Secretion](#), [Carbohydrate Homeostasis](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 μ g/ μ L

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months