

Datasheet for ABIN1409069

anti-NMUR2 antibody (AA 181-230) (Cy5.5)



()	ve	rvi	6	W
\sim	v C	1 V I	\sim	v v

100 μL
NMUR2
AA 181-230
Human
Rabbit
Polyclonal
This NMUR2 antibody is conjugated to Cy5.5
Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
KLH conjugated synthetic peptide derived from human NMUR2
IgG
IgG Human,Cow,Sheep,Pig,Horse,Rabbit
Human,Cow,Sheep,Pig,Horse,Rabbit
Human,Cow,Sheep,Pig,Horse,Rabbit
Human,Cow,Sheep,Pig,Horse,Rabbit Purified by Protein A.

coupled receptor FM-4, G-protein coupled receptor TGR-1, Neuromedin U receptor 2, Neuromedin-U receptor 2, NMU R2, NMU-R2, NMU-R2, NMUR2, NMUR2_HUMAN, TGR1

Background: Neuromedin U is a neuropeptide with high activity on smooth muscle. It is widely expressed in gastrointestinal systems and central nervous system (CNS). Peripheral activities of neuromedin U include smooth muscle stimulation, ion transport alterations in the gut and the regulation of local blood flow and adrenocortical function. Neuromedin U receptors 1 and 2 (NMUR1 and NMUR2) are multi-pass membrane proteins that belong to the G-protein coupled receptor 1 family of proteins. Both NMUR1 and NMUR2 act as receptors for the neuromedin U neuropeptide. NMUR1 is detected in peripheral organs, particularly in urogenital and gastrointestinal systems, with highest levels in testis. It?s expression in CNS is low, but the protein has been detected in cerebellum, hippocampus, dorsal root ganglia and spinal cord. NMUR2 is predominantly detected in central nervous system with highest levels detected in medulla oblongata, spinal cord and thalamus. It may also be detected in testis but has low levels of expression in peripheral tissues.

Gene ID:	56923
UniProt:	Q9GZQ4
Pathways:	Feeding Behaviour

Application Details

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months