

Datasheet for ABIN1409463

anti-TOR1A antibody (AA 56-130) (Cy3)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	TOR1A
Binding Specificity:	AA 56-130
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TOR1A antibody is conjugated to Cy3
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Torsin A
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig
Purification:	Purified by Protein A.

Target Details

Target:	TOR1A
Alternative Name:	Torsin A (TOR1A Products)

Target Details

Background:	Synonyms: DQ2, Dystonia 1, Dystonia 1 protein, Dyt1, Tor1a, Torsin A, Torsin family 1 member A, TOR1A_HUMAN. Background: A mutation of the DYT1 gene, which codes for TorsinA, has been identified as the cause of one form of autosomal dominantly inherited dystonia. Early-onset torsion dystonia is a movement disorder, characterized by twisting muscle contractures, that begins in childhood. Symptoms are believed to result from altered neuronal communication in the basal ganglia. TorsinA comprises 332 amino acids. TorsinA is widely expressed throughout the mouse central nervous system and is detected in the majority of neurons in nearly all regions. The proteins display cytoplasmic distribution, although in some types of neurons localization is perinuclear. TorsinA often performs chaperone-like functions that assist in the assembly, operation, or disassembly of protein complexes. The gene which encodes TorsinA has high homology to three additional mammalian genes and a nematode gene and distal similarity to the family of heat-shock proteins and the Clp protease family. The gene which encodes TorsinA maps to human chromosome 9q34.
Pathways:	SARS-CoV-2 Protein Interactome

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months