

## Datasheet for ABIN1409528 anti-CADPS2 antibody (AA 435-510) (HRP)



Go to Product page

| _ |   |   |    |    |   |
|---|---|---|----|----|---|
|   | W | 0 | rv | 10 | W |

| Quantity:  | 100 μL   |  |
|--|--|--|
| Target:  | CADPS2   |  |
| Binding Specificity:   | AA 435-510   |  |
| Reactivity:  | Human  |  |
| Host:  | Rabbit   |  |
| Clonality:   | Polyclonal   |  |
| Conjugate:   | This CADPS2 antibody is conjugated to HRP                          |  |
| Application:   | ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro))          |  |
|  |  |  |
| Product Details  |  |  |
| Product Details  Immunogen:  | KLH conjugated synthetic peptide derived from human CADPS2         |  |
|  | KLH conjugated synthetic peptide derived from human CADPS2         |  |
| Immunogen:   |  |  |
| Immunogen: Isotype:  | IgG  |  |
| Immunogen: Isotype: Predicted Reactivity:                              | lgG<br>Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse                     |  |
| Immunogen: Isotype: Predicted Reactivity: Purification:                | lgG<br>Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse                     |  |
| Immunogen: Isotype: Predicted Reactivity: Purification: Target Details | lgG Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse Purified by Protein A. |  |

Background: Calcium-dependent secretion activators (CAPS-1 and CAPS-2) are calcium-binding proteins that direct neurotransmitter and neuropeptide-filled vesicles to the cell membrane for secretory granule exocytosis. Both CAPS-1 and CAPS-2 are expressed primarily in the brain where they regulate the secretion of various substances. The CAPS proteins contain a PH domain that is essential for regulation of exocytosis, as well as regulation of phospholipid binding. Through their regulation of neurotrophin release from granule cells, CAPS proteins help to regulate cell fate during neuronal development. CAPS-1 is thought to regulate catecholamine release from neuronal cells, while CAPS-2 is thought to regulate release of both brain-derived neurotrophic factor and neurotrophin-3 from granule cells. Defects in the genes encoding CAPS-1 and CAPS-2 are implicated in impaired cerebral development and autism.

Pathways:

Synaptic Vesicle Exocytosis

## **Application Details**

Application Notes: WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions: For Research Use only

## Handling

| Format:            | Liquid   |  |
|--------------------|--|--|
| Concentration:     | 1 μg/μL  |  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |  |
| Preservative:      | ProClin  |  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |  |
| Handling Advice:   | Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.               |  |
| Storage:           | -20 °C   |  |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |  |
| Expiry Date:       | 12 months  |  |