

Datasheet for ABIN1409620 **anti-FADS2 antibody (Cy5)**



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	FADS2
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FADS2 antibody is conjugated to Cy5
Application:	Western Blotting (WB)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FADS2
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	FADS2
Alternative Name:	Fads2 (FADS2 Products)
Background:	Synonyms: D6D, Delta6 desaturase, Delta6 fatty acid desaturase, DES 6, DES6, FADS 2, FADSD 6, FADSD6, Fatty acid desaturase 2, linoleoyl-CoA desaturase delta-6-desaturase like 2, LLCDL 2, LLCDL2, SLL0262, TU 13, TU13, FADS2_HUMAN. Background: Members of the fatty acid desaturase (FADS) family, including FADS1, FADS2 and

Target Details

FADS3, regulate the desaturation of fatty acids by introducing double bonds between defined carbons of fatty acyl chains, thereby playing an essential role in the lipid metabolic pathway. Members of this family share N-terminal cytochrome b5-like domains, C-terminal multiple membrane-spanning desaturase regions and 3 histidine box motifs. FADS2 (fatty acid desaturase 2), also known as D6D, DES6, LLCDL2 or TU13, is a 444 amino acid multi-pass membrane protein that localizes to the endoplasmic reticulum and contains one cytochrome b5 heme-binding domain. Expressed in adult and fetal heart and in adult liver, brain, lung and retina, FADS2 functions as a component of a lipid metabolic pathway and catalyzes the first step in the pathway, namely the formation of unsaturated fatty acids from polyunsaturated fatty acids. Defects in the gene encoding FADS2 are the cause of cause of fatty acid delta-6-desaturase deficiency, an affliction that is characterized by skin abnormalities, corneal ulceration and growth failure. Multiple isoforms of FADS2 exist due to alternative splicing events.

Application Details

Application Notes:	IF(IHC-P)(1:50-200)
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months