

Datasheet for ABIN1409672 anti-NHLH2 antibody (AA 1-50) (HRP)



Go to Product page

| Overview | |
|-----------------------|---|
| Quantity: | 100 μL |
| Target: | NHLH2 |
| Binding Specificity: | AA 1-50 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This NHLH2 antibody is conjugated to HRP |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |
| Product Details | |
| Immunogen: | KLH conjugated synthetic peptide derived from human NHLH2 |
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Predicted Reactivity: | Mouse,Rat,Dog,Cow,Pig |
| Purification: | Purified by Protein A. |
| Target Details | |
| Target: | NHLH2 |
| Alternative Name: | NHLH2 (NHLH2 Products) |

Target Details

| Ranko | round: |
|-------|----------|
| Dacke | n Ouriu. |

Synonyms: HEN2, Nescient helix loop helix 2, NSCL, NSCL2, HEN2_HUMAN.

Background: The helix-loop-helix (HLH) structures are known motifs commonly found in membrane-active and DNA-binding proteins. The helix-loop-helix proteins HEN1 and HEN2 are DNA-binding proteins that may be involved in cell-type determination in the early nervous system. Studies of expression in normal tissues have demonstrated expression of NHLH1/NSCL-1 and NHLH2/NSCL-2, the genes encoding HEN1 and HEN2, in the developing central and peripheral nervous system, specifically in developing neurons.

Application Details

Application Notes:

WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Handling Advice: | Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |