

Datasheet for ABIN1410062

anti-Homeobox C5 antibody (AA 151-222) (HRP)



Go to Product page

()	ve	r\/i	Δ	۱۸/
\circ	V C	1 V		v v

Quantity:	100 μL	
Target:	Homeobox C5 (HOXC5)	
Binding Specificity:	AA 151-222	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Homeobox C5 antibody is conjugated to HRP	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)),	
	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human HOXC5	
	New York and The Control of the Cont	
Isotype:	IgG	
Isotype: Cross-Reactivity:		
	IgG	
Cross-Reactivity:	IgG Human, Mouse	
Cross-Reactivity: Predicted Reactivity:	IgG Human, Mouse Rat,Dog,Cow,Sheep	
Cross-Reactivity: Predicted Reactivity: Purification:	IgG Human, Mouse Rat,Dog,Cow,Sheep	

Target Details

Background:

Synonyms: CP11, homeo box C5, homeobox C5, Homeobox protein CP11, homeobox protein Hox C5, Homeobox protein Hox-3D, Homeobox protein Hox-C5, Hox 3D, HOX3, HOX3D, HOXC5, HXC5_HUMAN, OTTHUMP00000217386.

Background: The Hox homeobox genes encode proteins that are transcriptional regulators with an established role in embryonic development. HoxC5 is a 222 amino acid protein that localizes to the nucleus and contains one homeobox DNA-binding domain. Existing as multiple alternatively spliced isoforms, HoxC5 functions as a sequence-specific DNA-binding transcription factor that is part of a regulatory mechanism that provides cells with positional identities during development. The gene encoding HoxC5 maps to a cluster of Hox proteins on chromosome 12 that are essential for morphogenesis. Encoding over 1,100 genes within 132 million bases, chromosome 12 makes up about 4.5 % of the human genome. A number of skeletal deformities are linked to chromosome 12, including hypochondrogenesis, achondrogenesis and Kniest dysplasia.

Application Details

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

· · · · · · · · · · · · · · · · · · ·		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	

ш	Jand	lina
г	land	1111()
•	10110	9

Expiry Date:

12 months