

Datasheet for ABIN1410167

anti-TSKU antibody (AA 231-300) (Cy5.5)



Overview

Overview		
Quantity:	100 μL	
Target:	TSKU	
Binding Specificity:	AA 231-300	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TSKU antibody is conjugated to Cy5.5	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human Tsukushin/LRRC54	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Dog,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	
Target Details		
Target:	TSKU	
Alternative Name:	TSK/Tsukushin/LRRC54 (TSKU Products)	
Background:	Synonyms: E2 induced gene 4 protein, E2IG4, Leucine rich repeat containing protein 54,	

LRRC54, TSK, Tsukushi, TSK_HUMAN, TSKU.

Background: The leucine-rich (LRR) repeat is a 20-30 amino acid motif that forms a hydrophobic / horseshoe fold, allowing it to accommodate several leucine residues within a tightly packed core. All LRR repeats contain a variable segment and a highly conserved segment, the latter of which accounts for 11 or 12 residues of the entire LRR motif. LRRC54 (leucine-rich repeat-containing protein 54), also known as tsukushin, TSKU or E2-induced gene 4 protein (E2IG4), is a 353 amino acid secreted protein that likely localizes to the cell membrane and extracellular compartments. Involved in extracellular secretion and intracellular transport, LRRC54 can be induced by 17-beta-estradiol. Containing nine LRR repeat and a cleavable signal peptide, the gene encoding LRRC54 maps to human chromosome 11, which houses over 1,400 genes and comprises nearly 4 % of the human genome. Jervell and Lange-Nielsen syndrome, Jacobsen syndrome, Niemann-Pick disease, hereditary angioedema and Smith-Lemli-Opitz syndrome are associated with defects in genes that maps to chromosome 11.

Gene ID: 25987

UniProt: Q8WUA8

Application Details

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

	1	l:
-	ココロロ	urnen
	Hand	III IU

Expiry Date:

12 months