antibodies -online.com





anti-TAS2R7 antibody (AA 121-220) (Cy5)



Go to Product page

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	TAS2R7
Binding Specificity:	AA 121-220
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TAS2R7 antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human TAS2R7/T2R7	
Isotype:	IgG	
Predicted Reactivity:	Human	
Purification:	Purified by Protein A.	

Target Details

Target:	TAS2R7	
Alternative Name:	TAS2R7/T2R7 (TAS2R7 Products)	
Background:	Synonyms: STC7 4, T2R30, T2R6, T2R7, TA2R7_HUMAN, Tas2r130, Tas2r6, TAS2R7, Taste	

receptor family B member 4, Taste receptor type 2 member 7, TRB4.

Background: The sense of taste provides animals with valuable information about the quality and nutritional value of food. A family of G protein-coupled receptors are involved in taste perception and include T1R, which is involved in sweet and umami taste perception, and T2R, which is involved in bitter taste perception. Both types of taste receptors couple to various G proteins to initiate signal transduction cascades. Single taste receptor cells express a variety of T2Rs, suggesting that each cell is capable of recognizing multiple tastants. T2R6 (also designatedT2R30, mt2r42, STC 7-4 or taste receptor, type 2, member 130) is an integral membrane receptor protein in mice that may play a role in the perception of bitterness and in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate Alpha-gustducin, mediate PLC-Beta-2 activation and lead to the gating of TRPM5. T2R6 is expressed in subsets of taste receptor cells of the tongue and palate epithelium and exclusively in gustducin-positive cells. The human homolog of T2R6, designated T2R7 (TAS2R7, TRB4 or taste receptor, type 2, member 7) is a G protein-coupled receptor expressed in taste receptor cells of the tongue and palate epithelia.

Application Details

Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	

		1.	
\vdash	land	lın	
	ıaııu	1111	U

Expiry Date:

12 months