

# Datasheet for ABIN1410796 anti-CLN5 antibody (AA 61-120) (Cy5)



| Overview              |  |
|-----------------------|--|
| Quantity:             | 100 μL   |
| Target:               | CLN5   |
| Binding Specificity:  | AA 61-120  |
| Reactivity:           | Human  |
| Host:                 | Rabbit   |
| Clonality:            | Polyclonal   |
| Conjugate:            | This CLN5 antibody is conjugated to Cy5  |
| Application:          | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |
| Product Details       |  |
| Immunogen:            | KLH conjugated synthetic peptide derived from human CLN5   |
| Isotype:              | IgG  |
| Cross-Reactivity:     | Human  |
| Predicted Reactivity: | Mouse,Rat,Cow,Sheep,Pig,Horse,Rabbit   |
| Purification:         | Purified by Protein A.   |
| Target Details        |  |
| Target:               | CLN5   |
|                       |  |

### **Target Details**

#### Background:

Synonyms: Ceroid lipofuscinosis neuronal 5, Ceroid-lipofuscinosis neuronal protein 5, CLN5, CLN5\_HUMAN, NCL, Protein CLN5.

Background: Neuronal ceroid-lipofuscinose (NCL), also designated Batten disease, comprises a group of recessively inherited, progressive neurodegenerative diseases found in children. NCL is characterized by atrophy of the brain and an accumulation of lysosome derived fluorescent bodies found in many cells, especially neurons. Symptoms of NCL include a failure of psychomotor development, seizures, impaired vision and premature death. The eight genes/proteins associated with NCL are designated CLN1-CLN8. Mutations in six of these genes results in a distinct type of NCL-disease, the six genes/proteins are CLN1 (encoding PPT1, a protein thiolesterase), CLN2 (encodeing the serine protease TPP1), CLN3, CLN5, CLN6 and CLN8. A single base duplication mutation in dog and cow CLN5 has been shown to cause NCL.

# **Application Details**

| ۸  | -1::      | NI-+   |
|----|-----------|--------|
| ΑD | olication | notes. |

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

## Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |