

# Datasheet for ABIN1411068

# anti-SULT4A1 antibody (AA 221-284) (Cy7)



_					
	1//	r	Vİ	$\triangle$	۸/
	V		VI		/ V

Quantity:	100 μL	
Target:	SULT4A1	
Binding Specificity:	AA 221-284	
Reactivity:	Human, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SULT4A1 antibody is conjugated to Cy7	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human SULT4A1	
Isotype:	IgG	
Cross-Reactivity:	Human, Rat	
Predicted Reactivity:	Mouse,Dog,Cow,Sheep,Pig,Horse,Rabbit	

## **Target Details**

Purification:

Target: SULT4A1 Alternative Name: SULT4A1 (SULT4A1 Products)

Purified by Protein A.

### **Target Details**

#### Background:

Synonyms: BR STL 1, Brain sulfotransferase like protein, Brain sulphotransferase like, BRSTL1, hBR STL 1, hBR STL, Nervous system cytosolic sulfotransferase, Nervous system sulfotransferase, NST, Sulfotransferase 4A1, Sulfotransferase family 4A member 1, Sulfotransferase related protein, SULTX3, ST4A1\_HUMAN.

Background: Sulfation is an essential conjugation reaction that increases the water solubility of many compounds, thereby influencing their renal excretion and also resulting in the formation of active metabolites. SULT4A1 (sulfotransferase family 4A, member 1), whose alternative names include brain sulfotransferase-like protein, nervous system sulfotransferase, NST, SULTX3, hBR-STL-1, BRSTL1, BR-STL-1, MGC40032 and DJ388M5.3, is a 284 amino acid protein showing cytoplasmic localization. As a member of the sulfotransferase 1 family, SULT4A1 plays a role in elimination of xenobiotics, activation of procarcinogens and regulation of hormones. SULT4A1 is highly expressed in cerebral cortex and frontal lobe, with lower expression in cerebellum, temporal and occipital lobes. Two SULT4A1 isoforms exist to alternative splicing events. The gene encoding SULT4A1 maps to human chromosome 22q13.2, a region which has been implicated in predisposition to schizophrenia.

### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

ш	Jand	lina
г	land	1111()
•	10110	9

Expiry Date:

12 months