

[Go to Product page](#)

Datasheet for ABIN1411425

anti-CSDC2 antibody (AA 21-120) (Cy3)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | CSDC2 |
| Binding Specificity: | AA 21-120 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This CSDC2 antibody is conjugated to Cy3 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human PIPPIN |
| Isotype: | IgG |
| Predicted Reactivity: | Human, Mouse, Rat, Cow, Sheep, Pig, Horse, Chicken, Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|---|
| Target: | CSDC2 |
| Alternative Name: | CSDC2/PIPPIN (CSDC2 Products) |
| Background: | Synonyms: Cold shock domain containing C2 RNA binding, Cold shock domain containing |

Target Details

protein C2, Cold shock domain-containing protein C2, CSDC 2, Csdc2, CSDC2_HUMAN, dJ347H13.2, PIPPIN, RNA binding protein PIPPIn, RNA-binding protein PIPPIn.

Background: PIPPIN is a Y-box protein (also called cold-shock (CSD) domain-containing protein) and belongs to a family of highly conserved RNA-binding transcriptional regulators.

Predominantly expressed in brain cells and localizing to the nucleus and the cytoplasm, PIPPIN contains two RNA-binding motifs, namely PIP1 and PIP2, and one CSD domain. PIPPIN functions as a nucleic acid binding regulatory factor and is believed to participate in brain maturation. More specifically, PIPPIN binds to the 3'-UTR ends of the mRNAs encoding Histone H1 and Histone H3.3. This interaction requires all of the PIPPIN domains to work in concert as one functional protein. In addition, PIPPIN can be sumoylated in a thyroid hormone (T3)-dependent manner. This suggests that PIPPIN modification in response to extracellular stimuli may modulate the regulation of protein synthesis.

Application Details

| | |
|--------------------|--|
| Application Notes: | IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |

Handling

| | |
|--------------------|--|
| Format: | Liquid |
| Concentration: | 1 µg/µL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |