

Datasheet for ABIN1411496
anti-DYRK1A antibody (AA 81-170) (HRP)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	DYRK1A
Binding Specificity:	AA 81-170
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DYRK1A antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DYRK1A
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	DYRK1A
Alternative Name:	DYRK1A (DYRK1A Products)
Background:	Synonyms: Dual specificity tyrosine phosphorylation regulated kinase 1A, DYRK 1, DYRK 1A,

Target Details

DYRK , DYRK1 , DYRKA, HP 86, HP86, Minibrain Drosophila homolog, Minibrain homolog, MNB , MNB/DYRK protein kinase antibody, MNBH, Protein kinase minibrain homolog, Serine/threonine kinase MNB, DYR1A_HUMAN.

Background: Dyrk (for dual specificity tyrosine phosphorylation regulated kinase) is the homolog of the Drosophila mnbr (minibrain) gene which is required for neurogenesis. Dyrk is a dual-specificity tyrosine kinase and serine/threonine kinase, which is itself regulated by tyrosine phosphorylation. Several mammalian Dyrk related proteins have been identified and are thought to compose a family of dual specificity protein kinases. Dyrk family members, including Dyrk1A (dual specificity tyrosine-phosphorylation-regulated kinase 1A), Dyrk1B, Dyrk1C, Dyrk2, Dyrk3, Dyrk4A and Dyrk4B, are thought to be involved in diverse cellular functions. Localized to the nucleus and highly expressed in testis, muscle and the developing nervous system, Dyrk1A, also known as MNB or MNBH, functions to phosphorylate serine, threonine and tyrosine residues on various substrates involved in signaling pathways that regulate cell proliferation. Dyrk1A is a candidate gene for learning defects that are involved in Down's syndrome (DS), suggesting a possible role for Dyrk1A in the development of DS. Four isoforms of Dyrk1A exist due to alternative splicing events.

Pathways: [Mitotic G1-G1/S Phases](#)

Application Details

Application Notes: WB 1:300-5000
IHC-P 1:200-400
IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months