

Datasheet for ABIN1411556

anti-ARHGEF25/GEFT antibody (AA 251-340) (HRP)



	Go to Froduct page

()	ve	r\/i	Δ	۱۸/
\circ	V C	1 V		v v

Quantity:	100 μL	
Target:	ARHGEF25/GEFT (ARHGEF25)	
Binding Specificity:	AA 251-340	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ARHGEF25/GEFT antibody is conjugated to HRP	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human GEFT	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human, Mouse, Cow, Sheep, Pig, Rabbit	
Purification:	Purified by Protein A.	
Target Details		
Target:	ARHGEF25/GEFT (ARHGEF25)	
Alternative Name:	GEFT (ARHGEF25 Products)	

Target Details

Background:

Synonyms: ARHGEF25, ARHGP_HUMAN, GEFT, p63RhoGEF, Guanine nucleotide exchange factor GEFT, p63RhoGEF, RAC/CDC42 exchange factor, Rac/Cdc42/Rho exchange factor GEFT, Rho guanine nucleotide exchange factor 25, RhoA/RAC/CDC42 exchange factor, RhoA/Rac/Cdc42 guanine nucleotide exchange factor GEFT.

Background: GEFT is a 580 amino acid cytoplasmic protein that is highly expressed in excitable tissues such as brain, heart and muscle, and weakly expressed in small intestine, colon, liver, placenta and lung. GEFT may play a role in actin cytoskeleton reorganization in different tissues since its activation induces formation of actin stress fibers. GEFT works as a guanine nucleotide exchange factor for the Rho family of small GTPases and links specifically to G alpha q/11-coupled receptors in Rho A activation. GEFT is an important regulator of processes involved in axon and dendrite formation. Involved in skeletal myogenesis, GEFT seems to be an exchange factor primarily for Rac 1 in neurons. Existing as two alternatively spliced variants, GEFT contains a DH (DBL-homology) domain and a PH domain.

Application Details

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Liquid
1 μg/μL
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
ProClin
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
-20 °C
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

	1	l:
-	ココロロ	urnen
	Hand	III IU

Expiry Date:

12 months