## antibodies -online.com

Datasheet for ABIN1411635





anti-ITM2C antibody (AA 3-88) (Cy3)

0	:
U J\/∠r	view
$\circ$	V I C V V

Quantity:	100 μL
Target:	ITM2C
Binding Specificity:	AA 3-88
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ITM2C antibody is conjugated to Cy3
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
	( a.a aa.a.a.a acadama) ( (p))

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human ITM2C
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

## **Target Details**

Target:	ITM2C
Alternative Name:	ITM2C (ITM2C Products)
Background:	Synonyms: BRI3, BRICD2C, BRICHOS domain containing 2C, Cerebral protein 14, CT-BRI3, E25,

Storage Comment:

12 months

Expiry Date:

Target Details	
	E25C , Integral membrane protein 2C, Integral membrane protein 3, Itm2c, ITM2C_HUMAN,
	ITM3 antibodyNPD018, Transmembrane protein BRI3.
	Background: Negative regulator of beta amyloid peptide production. May inhibit the processing
	of APP by blocking its access to alpha-and beta-secretase. Binding to the beta-secretase-
	cleaved APP C-terminal fragment is negligible, suggesting that ITM2C is a poor gamma-
	secretase cleavage inhibitor. May play a role in TNF-induced cell death and neuronal
	differentiation.
Gene ID:	81618
UniProt:	Q9NQX7
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.