

[Go to Product page](#)

Datasheet for ABIN1411662

anti-Shootin-1 (SHTN1) (AA 21-130) antibody (Cy7)

Overview

Quantity:	100 µL
Target:	Shootin-1 (SHTN1)
Binding Specificity:	AA 21-130
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	Cy7
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Shootin/KIAA1598
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Cow,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	Shootin-1 (SHTN1)
Alternative Name:	Shootin/KIAA1598 (SHTN1 Products)

Target Details

Background:	<p>Synonyms: DKFZp686A0439, hypothetical protein FLJ11122, Kiaa1598, MGC40476, Putative uncharacterized protein KIAA1598, Shootin 1, Shootin-1, SHOT1_HUMAN.</p> <p>Background: Shootin1 is a 631 amino acid protein that belongs to the shootin family. The shootin1 protein contains three coiled-coil domains, a proline-rich region and interacts with RUFY3. Shootin1 is involved in the generation of internal asymmetric signals required for neuronal polarization. The shootin1 protein acts upstream of PI3K (phosphoinositide 3-kinase), by being required for spatially localized PI3K activity. By accumulating asymmetrically in a single neurite before polarization, shootin1 leads to axon induction for polarization, additionally the absence of shootin1 from the nascent axon's siblings by competition prevents the formation of surplus axons. Existing as seven alternatively spliced isoforms, the shootin1 gene is conserved in chimpanzee, dog, mouse, rat, chicken and zebrafish, and maps to human chromosome 10q25.3.</p>
-------------	---

Application Details

Application Notes:	<p>IF(IHC-P) 1:50-200</p> <p>IF(IHC-F) 1:50-200</p> <p>IF(ICC) 1:50-200</p>
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months