antibodies -online.com





anti-LGI3 antibody (AA 101-220) (Cy5)



	do to i roddot page

()	1 /	0	rv	/ 1 /	71	Α.
	1//	\vdash	1 \/	16		1/1/
\sim	v	\sim	1 V	١,	_	v v

Quantity:	100 μL
Target:	LGI3
Binding Specificity:	AA 101-220
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LGI3 antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human LGI3
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Sheep,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	LGI3
Alternative Name:	LGI3 (LGI3 Products)

Target Details

Background:

Synonyms: Leucine rich repeat LGI family member 3; Leucine-rich glioma inactivated protein 3; Leucine-rich glioma-inactivated protein 3; Leucine-rich repeat LGI family member 3; LGI1 like protein 4; LGI3; LGI3_HUMAN; LGIL4.

Background: The leucine-rich (LRR) repeat is a 20-30 amino acid motif that forms a hydrophobic å/ horseshoe fold, allowing it to accommodate several leucine residues within a tightly packed core. All LRR repeats contain a variable segment and a highly conserved segment, the latter of which accounts for 11 or 12 residues of the entire LRR motif. Leucine-rich glioma-inactivated protein 3 (LGI3), also known as LGI1-like protein 4 (LGIL4) or leucine-rich repeat LGI family member 3, is a 548 amino acid secretory protein. LGI3 contains five LRR repeats and seven EAR repeats. Widely expressed, with highest levels in brain and lung, LGI3 has been shown to colocalize with endocytosis-associated proteins, lipid raft markers, and Syntaxin. The gene encoding LGI3 maps to chromosome 8p21.3.

Application Details

Δn	plication	Notes:
\neg	piication	INOICS.

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months