

Datasheet for ABIN1411781

anti-MAP3K4 antibody (AA 1351-1450) (Cy5.5)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	MAP3K4
Binding Specificity:	AA 1351-1450
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MAP3K4 antibody is conjugated to Cy5.5
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MAP3K4/MEK kinase-4
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	MAP3K4
Alternative Name:	MAP3K4/MEK kinase-4 (MAP3K4 Products)
Background:	Synonyms: FLJ42439, KIAA0213, M3K4_HUMAN, MAP three kinase 1, MAP3K4, MAPK/ERK

Target Details

kinase kinase 4, MAPKKK4 antibodyMEK kinase 4, MEKK 4, MEKK4, Mitogen-activated protein kinase kinase kinase 4, MTK1, PRO0412, RP3-473J16.4

Background: Mitogen-activated protein (MAP) kinase cascades are activated by various extracellular stimuli including growth factors. The MEK kinases (also designated MAP kinase kinase kinases, MKKKs, MAP3Ks or MEKKs) phosphorylate and thereby activate the MEKs (also called MAP kinase kinases or MKKs), including ERK, JNK and p38. These activated MEKs in turn phosphorylate and activate the MAP kinases. The MEK kinases include Raf-1, Raf-B, Mos, MEK kinase-1, MEK kinase-2, MEK kinase-3, MEK kinase-4, ASK 1 (MEK kinase-5) and MAP3K6 (MEK kinase-6). MEK kinase-1 has been shown to phosphorylate MEK-1 via Raf-independent pathway. Evidence suggests that MEK-3 is preferentially activated by MEK kinase-3 and that MEK-4 is activated by both MEK kinase-2 and MEK kinase-3. MEK kinase-4 has been shown to specifically activate the JNK path-way. ASK1 activates both MEK-4 and MEK-3/MEK-6 pathways.

Pathways: [MAPK Signaling](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months