antibodies -online.com







anti-SLITRK2 antibody (AA 311-410) (Cy5)



\sim	4.4	D		page
1-()	17	$\mathbf{P}_{\mathbf{r}}$	וי או וו	nana
\cup	w	1 100	IUCL	Dauc

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	SLITRK2
Binding Specificity:	AA 311-410
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SLITRK2 antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SLITRK2
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit,Guinea Pig
Purification:	Purified by Protein A.

Target Details

Target:	SLITRK2
Alternative Name:	SLITRK2 (SLITRK2 Products)
Background: Synonyms: CXorf2, KIAA1854, SLIK2_HUMAN, SLIT and NTRK like protein 2, SLIT and N	

like protein 2, SLITL1, SLITRK 2, Slitrk2.

Background: SLITRK family proteins are integral membrane proteins that have a C-terminal domain that is partially similar to TRK neurotrophin receptor proteins and two leucine-rich repeat (LRR) domains that are similar to those of SLIT proteins. SLITRK2 (SLIT and NTRK-like protein 2) is a 845 amino acid single-pass type I membrane protein that contains 14 LRR (leucine-rich) repeats and is expressed in neural tissues, with highest levels found in adult cerebral cortex. Overexpression of SLITRK2 leads to inhibition of unipolar neurites in cultured cells, suggesting that it suppresses neurite outgrowth. Inhibitory activity of SLITRK2 is localized to its C-terminal intracellular domain and without this region the protein induces neurite outgrowth. Variants in the gene encoding SLITRK2 may contribute to the development of bipolar disorder, autism spectrum disorder and schizophrenia. There are two isoforms of SLITRK2 that are produced as a result of alternative splicing events.

Application Details

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months