Datasheet for ABIN1412204 anti-SLITRK3 antibody (AA 61-160) (HRP)

antibodies.com



$\sim$							
( )	۱/	0	r\	/1	$\cap$	۱۸	1
Ο	v	C	1 \	/ I	C	V١	/

Quantity:	100 µL	
Target:	SLITRK3	
Binding Specificity:	AA 61-160	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SLITRK3 antibody is conjugated to HRP	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SLITRK3
Isotype:	lgG
Predicted Reactivity:	Human,Mouse,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	SLITRK3
Alternative Name:	SLITRK3 (SLITRK3 Products)
Background: Synonyms: KIAA0848; SLIT and NTRK like protein 3; SLITRK 3; SLITRK 3; SLIK3_HU	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1412204 | 03/07/2024 | Copyright antibodies-online. All rights reserved. Background: The leucine-rich (LRR) repeat is a 20-30 amino acid motif that forms a hydrophobic å/ horseshoe fold, allowing it to accommodate several leucine residues within a tightly packed core. All LRR repeats contain a variable segment and a highly conserved segment, the latter of which accounts for 11 or 12 residues of the entire LRR motif. SLITRK3 (SLIT and NTRK-like family, member 3) is a 977 amino acid single-pass type I membrane protein that contains 20 LRR repeats and belongs to the SLITRK family. Expressed at highest levels in cerebral cortex, SLITRK3 is also found in adult and fetal neural tissues and some astrocytic brain tumors. SLITRK3 functions to suppress neurite outgrowth and plays a role in the regulation of neuronal function. SLITRK3 is encoded by a gene that maps to human chromosome 3, which houses over 1,100 genes, including a chemokine receptor (CKR) gene cluster and a variety of human cancer-related gene loci.

## Application Details

1 1		
Application Notes:	WB 1:300-5000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN1412204 | 03/07/2024 | Copyright antibodies-online. All rights reserved.