

[Go to Product page](#)

Datasheet for ABIN1412386  
**anti-NOS1AP antibody (AA 128-170) (Cy5)**

## Overview

Quantity:	100 µL
Target:	NOS1AP
Binding Specificity:	AA 128-170
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NOS1AP antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CAPON
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Horse
Purification:	Purified by Protein A.

## Target Details

Target:	NOS1AP
Alternative Name:	CAPON ( <a href="#">NOS1AP Products</a> )
Background:	Synonyms: NOS1AP, C terminal PDZ domain ligand of neuronal nitric oxide synthase CAPON, C

## Target Details

terminal PDZ domain ligand of neuronal nitric oxide synthase, C terminal PDZ ligand of neuronal nitric oxide synthase protein, C-terminal PDZ ligand of neuronal nitric oxide synthase protein, CAPON, CAPON\_HUMAN, Carboxyl terminal PDZ ligand of neuronal nitric oxide synthase protein, Carboxyl-terminal PDZ ligand of neuronal nitric oxide synthase protein, Ligand of neuronal nitric oxide synthase with carboxyl terminal PDZ domain, MGC138500, Nitric oxide synthase 1 neuronal adaptor protein, Nitric oxide synthase 1 adaptor protein.

Background: CAPON (carboxy-terminal PDZ ligand of nNOS) selectively binds within the 100 amino acid PDZ domain of the neuronal nitric oxide synthase (nNOS), but not to endothelial NOS or inducible NOS, and sequesters nNOS in the cytosol. Biosynthesis of the neurotransmitter nitric oxide (NO) requires the association of nNOS with various synaptic proteins, including syntrophin, postsynaptic density (PSD)95 and PSD93 through a scaffolding PDZ domain. These proteins facilitate the transport of nNOS to the plasma membrane, where it is catalytically activated by NMDA-receptor mediated calcium channels. The association of nNOS with PSD95 or PSD93 is regulated by CAPON. The carboxy terminus of CAPON binds to the PDZ domain, competes with PSD95 and PSD93 for binding to nNOS and in turn prevents the translocation and catalytic activation of nNOS.

## Application Details

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

## Handling

---

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

---

Expiry Date: 12 months