



[Go to Product page](#)

Datasheet for ABIN1412601

## anti-Trace Amine Associated Receptor 8 (TAAR8) (AA 51-150) antibody (Cy3)

### Overview

Quantity:	100 µL
Target:	Trace Amine Associated Receptor 8 (TAAR8)
Binding Specificity:	AA 51-150
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	Cy3
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human G protein coupled receptor 102
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Cow, Sheep, Pig, Rabbit
Purification:	Purified by Protein A.

### Target Details

Target:	Trace Amine Associated Receptor 8 (TAAR8)
Alternative Name:	GPR102/TAAR8 ( <a href="#">TAAR8 Products</a> )
Background:	Synonyms: G protein coupled receptor 102, G-protein coupled receptor 102, GPR102, TA5,

## Target Details

---

TAAR8, TAAR8\_HUMAN, TaR-5, TaR-8, TAR5, Trace amine associated receptor 8, Trace amine receptor 5, Trace amine receptor 8, Trace amine-associated receptor 8, TRAR5, GPCR GPR102. Background: Trace amines are endogenous molecules structurally related to classical biogenic amines that are linked to psychiatric conditions. A family of G-protein coupled receptors referred to as trace-amine-associated receptors (TAAR) are activated by trace amines and are present in very low levels in mammalian tissue. TaRs contain several structural features that are similar to the rhodopsin -adrenergic receptor superfamily, including the positions of the seven transmembrane regions that provide common ligand-binding pockets as well as the short N- and C-terminal domains. TAAR proteins are potential targets for drugs of abuse, such as amphetamine and MDMA, as well as neuropsychiatric disorders including schizophrenia, depression, and attention deficit disorder.

## Application Details

---

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months