

Datasheet for ABIN1412726 anti-UNCX antibody (AA 145-150) (HRP)



Go to Product page

Overview	
Quantity:	100 μL
Target:	UNCX
Binding Specificity:	AA 145-150
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UNCX antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human UNCX
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Pig,Chicken
Purification:	Purified by Protein A.
Target Details	
Target:	UNCX
Alternative Name:	UNCX (UNCX Products)

Target Details

Background:

Synonyms: Homeobox protein unc-4 homolog, Homeobox protein Uncx4.1, tcag7.1272, UNC homeobox, Unc4.1 homeobox, UNC4_HUMAN, Uncx, UNCX4.1.

Background: Members of the paired homeobox family play a role in regulating cell development and pattern formation during embryonic stages. UNCX (UNC homeobox), also known as UNCX4.1, is a 531 amino acid nuclear transcription factor involved in neurogenesis and somitogenesis. Containing one homeobox DNA-binding domain, UNCX belongs to the paired homeobox family and UNC4 subfamily. UNCX assists in the formation of connections between hypothalamic neurons and the pituitary, which is necessary for central neurons to deliver hormones into peripheral blood. UNCX also plays a role in maintaining differentiation of the axial skeleton and acts upstream of Pax-9. The gene encoding UNCX maps to human chromosome 7, which houses over 1,000 genes and comprises nearly 5 % of the human genome. Chromosome 7 has been linked to Osteogenesis imperfecta, Pendred syndrome, Lissencephaly, Citrullinemia and Shwachman-Diamond syndrome.

Application Details

Restrictions:

Application Notes:	WB 1:300-5000

IHC-P 1:200-400 IHC-F 1:100-500

For Research Use only

Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

	1	l:
-	ココロロ	urnen
	Hand	III IU

Expiry Date:

12 months