

## Datasheet for ABIN1413252

## anti-PNCK antibody (AA 31-160) (Cy7)



Go to Product page

_						
	V	$\triangle$	r۱	/1	$\triangle$	Λ/
	' V '		ΙV			v v

Quantity:	100 μL		
Target:	PNCK		
Binding Specificity:	AA 31-160		
Reactivity:	Human		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This PNCK antibody is conjugated to Cy7		
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))		
Product Details			
Immunogen:	KLH conjugated synthetic peptide derived from human PNCK		
Isotype:	IgG		
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Pig		
Purification:	Purified by Protein A.		
Target Details			
Target:	PNCK		
Alternative Name:	PNCK (PNCK Products)		
Background:	Synonyms: Calcium/calmodulin dependent protein kinase type 1B, Calcium/calmodulin-		

dependent protein kinase type 1B, CaM KI beta, CaM kinase I beta, CaM kinase IB, CaM-KI beta, CaMK1b, CaMKI beta, CaMKI-beta, EC 2.7.11.17, KCC1B\_HUMAN, MGC45419, PNCK, Pregnancy up regulated non ubiquitously expressed CaM kinase, Pregnancy up-regulated non-ubiquitously-expressed CaM kinase.

Background: The Ca2+/calmodulin-dependent protein kinases (CaMKs) comprise a structurally related subfamily of serine/threonine kinases. CaMKI Beta (Ca2+/calmodulin-dependent protein kinase type 1B), also known as PNCK (pregnancy up-regulated non-ubiquitously expressed CaM kinase) or BSTK3, is a 343 amino acid protein that localizes to both the nucleus and the cytoplasm and contains one protein kinase domain. Existing as multiple alternatively spliced isoforms, CaMKI Beta functions to catalyze the ATP-dependent phosphorylation of CaMKI, an event that activates CaMKI activity and may be important for Ca2+-triggered signaling cascades within the cell. The gene encoding CaMKI Beta maps to human chromosome X, which contains nearly 153 million base pairs and houses over 1,000 genes.

## **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months